Regularity of Wasserstein barycenters

Dissertation defense

Jianyu Ma

[supervisor]: JÉRÔME BERTRAND

October 16, 2025 Institut de Mathématiques de Toulouse

Barycenter (center of mass)

Barycenter z_{μ} averages points in a metric space (E,d) according to the probability measure μ on E, in the sense that

$$\int_E d(z_{\mu}, y)^2 d\mu(y) = \inf_{x \in E} \int_E d(x, y)^2 d\mu(y).$$

Wasserstein barycenters: averaging measures

- 1. $(E,d) \rightarrow (\mathcal{W}_2(E),d_W)$
- 2. $\mu \to \mathbb{P}$ and $z_{\mu} \to \mu_{\mathbb{P}}$
- $\mathcal{W}_2(E)$ is the set of probability measures on E with finite second-order moments, including $\mu_\mathbb{P}$

- Structure via base metric space
- Feature-preserving after averaging:e.g. Gaussians remain Gaussian

Barycenter (center of mass)

Barycenter z_{μ} averages points in a metric space (E,d) according to the probability measure μ on E, in the sense that

$$\int_E d(z_{\mu}, y)^2 d \mu(y) = \inf_{x \in E} \int_E d(x, y)^2 d \mu(y).$$

Wasserstein barycenters: averaging measures

- 1. $(E, d) \rightarrow (\mathcal{W}_2(E), d_W)$
- 2. $\mu \to \mathbb{P}$ and $z_{\mu} \to \mu_{\mathbb{P}}$

 $\mathcal{W}_2(E)$ is the set of probability measures on E with finite second-order moments, including $\mu_\mathbb{P}.$

- Structure via base metric space
- Feature-preserving after averaging:
 e.g. Gaussians remain Gaussian

Example: $\mu_{\mathbb{P}}$ of $\mathbb{P}:=rac{1}{2}\delta_{
u_1}+rac{1}{2}\delta_{
u_2}$

Consider the earth surface (E, d) with uniform measures ν_1, ν_2 supported on two regions.



$$+$$
 $\stackrel{\text{barycenter}}{\longrightarrow}$ \swarrow (11ama)

Wasserstein barycenters: averaging measures


- 1. $(E, d) \rightarrow (\mathcal{W}_2(E), d_W)$
- 2. $\mu \to \mathbb{P}$ and $z_{\mu} \to \mu_{\mathbb{P}}$

 $\mathcal{W}_2(E)$ is the set of probability measures on E with finite second-order moments, including $\mu_{\mathbb{P}}.$

- Structure via base metric space
- Feature-preserving after averaging: e.g. Gaussians remain Gaussian

Example: averaging Gaussian data

Two sensors returning estimates $\mathcal{N}(5,1)$ and $\mathcal{N}(9,0.8)$, then averaged in two different ways.



Wasserstein barycenters: averaging measures

- 1. $(E, d) \rightarrow (\mathcal{W}_2(E), d_W)$
- 2. $\mu \to \mathbb{P}$ and $z_{\mu} \to \mu_{\mathbb{P}}$

 $\mathcal{W}_2(E)$ is the set of probability measures on E with finite second-order moments, including $\mu_\mathbb{P}.$

- Structure via base metric space
- ► Feature-preserving after averaging: e.g. Gaussians remain Gaussian

Absolute continuity

Singularity



Example: coloring a cat with brush or pen

Gradient with brush

hue values change continuously \implies absolute continuity

Tattoo with pen

curves have zero area \implies singularity

Absolute continuity [manifolds]

Singularity

- [Contributions of the thesis]
 - 2 now class of displacement functions



Example: coloring a cat with brush or pen

Gradient with brush

hue values change continuously \implies absolute continuity

Tattoo with pen curves have zero area ⇒ singularity

Absolute continuity [manifolds]

Singularity

- [Contributions of the thesis]

 1 lower Ricci curvature bound suffices
 - 2 new class of displacement functiona



Example: coloring a cat with brush or pen

Gradient with brush

hue values change continuously \implies absolute continuity

Tattoo with pen curves have zero area ⇒ singularity

Absolute continuity [manifolds]

Singularity

[Contributions of the thesis]

- 1. lower Ricci curvature bound suffices
- 2. new class of displacement functional



Example: coloring a cat with brush or pen

Gradient with brush

hue values change continuously \implies absolute continuity

Tattoo with pen

curves have zero area \implies singularity


Absolute continuity [manifolds] [Contributions of the thesis]

- 1. lower Ricci curvature bound suffices
- 2. new class of displacement functional

Singularity [metric trees]

[Contributions of the thesis]

- reduction and localization techniques
- 4. rigid properties of barycenters on $\mathbb R$



Example: singularity at the branching point


$$\mathbb{P} := \frac{1}{3}\delta_{\nu_1} + \frac{1}{3}\delta_{\nu_2} + \frac{1}{3}\delta_{\nu_3} \implies \mu_{\mathbb{P}} = \delta_0$$

Absolute continuity [manifolds] [Contributions of the thesis]

- 1. lower Ricci curvature bound suffices
- 2. new class of displacement functional

Singularity [metric trees] [Contributions of the thesis]

- 3. reduction and localization techniques
- 4. rigid properties of barycenters on \mathbb{R}



Reduction: flatten w.r.t. a fixed edge \vec{e}

Localization: looping over all edges

Absolute continuity [manifolds] [Contributions of the thesis]


1. lower Ricci curvature bound suffices

2. new class of displacement functional

Singularity [metric trees]

[Contributions of the thesis]

- 3. reduction and localization techniques
- 4. rigid properties of barycenters on $\mathbb R$



Example: being singular is a rigid property

 $\mu_{\mathbb{P}} \in \mathcal{W}_2(\mathbb{R})$ is singular \implies \mathbb{P} -almost every u is singular

proper: bounded + closed => compact

Existence, uniqueness, and consistency

Let (E,d) be a proper metric space. Any $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(E))$ has a barycenter.

[F. Santambrogio, 2015] [Y.-H. Kim and B. Pass, 2017] Let (M, d_g) be a complete Riemannian manifold $\mathbb{P}(\text{a.c. measures}) > 0 \implies \mu_{\mathbb{P}}$ is unique.

[T. Le Goule and J.-M. Loubes, 2017] Let (E,d) be a proper metric space. If a sequence $\mathbb{P}_j \to \mathbb{P}$, then $\mu_{\mathbb{P}_j} \to \mu_{\mathbb{P}}$ up to a subsequence. a.c. = absolutely continuous

Absolute continuity

[M. Agueh and G. Carlier, 2011] Let $\nu_1, \ldots, \nu_n \in \mathcal{W}_2(\mathbb{R}^m)$ be n measures such that ν_1 is a.c. with bounded density. Then the unique barycenter $\mu_{\mathbb{P}}$ of $\mathbb{P} := \sum_{i=1}^n \lambda_i \, \delta_{\nu_i}$ is a.c. with

Let (M,d_0) be a compact Riemannian manifold. If a measure $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive massit to a set of a.c. measures with uniformly bounded density, then its unique barycenter $\mu_{\mathbb{P}}$ is a.c. with bounded density.

 ${\tt proper: bounded + closed} \implies {\tt compact}$

Existence, uniqueness, and consistency

[T. Le Gouic and J.-M. Loubes, 2017] Let (E, d) be a proper metric space. Any $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(E))$ has a barycenter.

[F. Santambrogio, 2015] [Y.-H. Kim and B. Pass, 2017] Let (M, d_g) be a complete Riemannian manifold $\mathbb{P}(\text{a.c. measures}) > 0 \implies \mu_{\mathbb{P}}$ is unique.

[T. Le Goule and J.-M. Loubes, 2017] Let (E,d) be a proper metric space. If a sequence $\mathbb{P}_{\!J} \to \mathbb{P}$, then $\mu_{\mathbb{P}_{\!J}} \to \mu_{\mathbb{P}}$ up to a subsequence. a.c. = absolutely continuous

Absolute continuity

M. Agueh and G. Carlier, 2011] Let $\nu_1, \ldots, \nu_n \in \mathcal{W}_2(\mathbb{R}^m)$ be n measures such that ν_1 is a.c. with bounded density. Then the unique parycenter $\mu_{\mathbb{P}}$ of $\mathbb{P} := \sum_{i=1}^n \lambda_i \, \delta_{\nu_i}$ is a.c. with bounded density.

Let (M,d_0) be a compact Riemannian manifold. If a measure $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive massive to a set of a.c. measures with uniformly bounded density, then its unique barycenter $\mu_{\mathbb{P}}$ is a.c. with bounded density.

 $\texttt{proper: bounded + closed} \implies \texttt{compact}$

Existence, uniqueness, and consistency

T. Le Gouic and J.-M. Loubes, 2017

Let (E,d) be a proper metric space. Any $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(E))$ has a barycenter.

 $[F.\ Santambrogio,\ 2015]$

[Y.-H. Kim and B. Pass, 2017]

Let (M, d_g) be a complete Riemannian manifold. $\mathbb{P}(\text{a.c. measures}) > 0 \implies \mu_{\mathbb{P}}$ is unique.

[T. Le Goule and J.-M. Loubes, 2017] Let (E,d) be a proper metric space. If a sequence $\mathbb{P}_j \to \mathbb{P}$, then $\mu_{\mathbb{P}_j} \to \mu_{\mathbb{P}}$ up to a subsequence. a.c. = absolutely continuous

Absolute continuity

[M. Agueh and G. Carlier, 2011]

Let $u_1,\dots,
u_n \in \mathcal{W}_2(\mathbb{R}^m)$ be n measures such that u_1 is a.c. with bounded density. Then the unique parycenter $u_\mathbb{P}$ of $\mathbb{P} := \sum_{i=1}^n \lambda_i \, \delta_{
u_i}$ is a.c. with bounded density.

[Y.-H. Kim and B. Pass, 2017] Let (M,d_g) be a compact Riemannian manifold. If a measure $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive massive to a set of a.c. measures with uniformly bounded density, then its unique barycenter $\mu_{\mathbb{P}}$ is a.c. with bounded density.

 $\texttt{proper: bounded + closed} \implies \texttt{compact}$

Existence, uniqueness, and consistency

T. Le Gouic and J.-M. Loubes, 2017

Let (E,d) be a proper metric space. Any $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(E))$ has a barycenter.

[F. Santambrogio, 2015]

[Y.-H. Kim and B. Pass, 2017]

Let $(M,d_{\rm g})$ be a complete Riemannian manifold. $\mathbb{P}({\rm a.c.\ measures})>0 \implies \mu_{\mathbb{P}}$ is unique.

[T. Le Gouic and J.-M. Loubes, 2017] Let (E,d) be a proper metric space. If a sequence $\mathbb{P}_j \to \mathbb{P}$, then $\mu_{\mathbb{P}_j} \to \mu_{\mathbb{P}}$ up to a subsequence.

a.c. = absolutely continuous

Absolute continuity

[M. Agueh and G. Carlier, 2011]

Let $u_1,\ldots, \nu_n \in \mathcal{W}_2(\mathbb{R}^m)$ be n measures such that u_1 is a.c. with bounded density. Then the unique barycenter $u_\mathbb{P}$ of $\mathbb{P} := \sum_{i=1}^n \lambda_i \, \delta_{\nu_i}$ is a.c. with bounded density.

[Y.-H. Kim and B. Pass, 2017] Let (M,d_{g}) be a compact Riemannian manifold. If a measure $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to a set of a.c. measures with uniformly bounded density, then its unique barycenter $\mu_{\mathbb{P}}$ is a.c. with bounded density.

proper: bounded + closed \implies compact

Existence, uniqueness, and consistency

T. Le Gouic and J.-M. Loubes, 2017

Let (E, d) be a proper metric space. Any $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(E))$ has a barycenter.

[F. Santambrogio, 2015]

[Y.-H. Kim and B. Pass, 2017]

Let $(M,d_{\rm g})$ be a complete Riemannian manifold. $\mathbb{P}({\rm a.c.\ measures})>0 \implies \mu_{\mathbb{P}}$ is unique.

[T. Le Gouic and J.-M. Loubes, 2017]

Let (E,d) be a proper metric space. If a sequence $\mathbb{P}_j \to \mathbb{P}$, then $\mu_{\mathbb{P}_j} \to \mu_{\mathbb{P}}$ up to a subsequence.

a.c. = absolutely continuous

Absolute continuity

[M. Agueh and G. Carlier, 2011]

Let $\nu_1,\dots,\nu_n\in\mathcal{W}_2(\mathbb{R}^m)$ be n measures such that ν_1 is a.c. with bounded density. Then the unique barycenter $\mu_\mathbb{P}$ of $\mathbb{P}:=\sum_{i=1}^n\lambda_i\,\delta_{\nu_i}$ is a.c. with bounded density.

Y.-H. Kim and B. Pass, 2017

Let (M,d_g) be a compact Riemannian manifold. If a measure $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to a set of a.c. measures with uniformly bounded density, then its unique barycenter $\mu_{\mathbb{P}}$ is a.c. with bounded density.

proper: bounded + closed \implies compact

Existence, uniqueness, and consistency

T. Le Gouic and J.-M. Loubes, 2017

Let (E, d) be a proper metric space. Any $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(E))$ has a barycenter.

[F. Santambrogio, 2015]

[Y.-H. Kim and B. Pass, 2017]

Let $(M,d_{\rm g})$ be a complete Riemannian manifold. $\mathbb{P}({\rm a.c.\ measures})>0 \implies \mu_{\mathbb{P}}$ is unique.

[T. Le Gouic and J.-M. Loubes, 2017]

Let (E,d) be a proper metric space. If a sequence $\mathbb{P}_j \to \mathbb{P}$, then $\mu_{\mathbb{P}_j} \to \mu_{\mathbb{P}}$ up to a subsequence.

a.c. = absolutely continuous

Absolute continuity

[M. Agueh and G. Carlier, 2011

Let $\nu_1,\dots,\nu_n\in\mathcal{W}_2(\mathbb{R}^m)$ be n measures such that ν_1 is a.c. with bounded density. Then the unique barycenter $\mu_\mathbb{P}$ of $\mathbb{P}:=\sum_{i=1}^n\lambda_i\,\delta_{\nu_i}$ is a.c. with bounded density.

[Y.-H. Kim and B. Pass, 2017]

Let (M,d_g) be a compact Riemannian manifold. If a measure $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to a set of a.c. measures with uniformly bounded density, then its unique barycenter $\mu_\mathbb{P}$ is a.c. with bounded density.

[J. Ma, 2023]

Let (M,d_g) be a complete Riemannian manifold with a lower Ricci curvature bound. If a measure $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to the set of a.c. probability measures, then its unique barycenter $\mu_{\mathbb{P}}$ is also a.c..

Improvements compared to the previous work
[Eliminate technical assumption]
a.c. with uniformly bounded density functions

[No obscure dependency on compactness] confirm the impact of lower Ricci curvature bounce on the regularity of barycenters

 $\mathbb{P} = \sum_{i=1}^n \lambda_i \, \delta_{
u_i} \implies \mu_{\mathbb{P}}$ can be constructed.

However, when \mathbb{P} is not finitely supported, its barycenter $\mu_{\mathbb{P}}$ is only accessible via consistency: as a limit of constructable barycenters $\mu_{\mathbb{P}_j}$ with \mathbb{P}_j finitely supported.

[Quantitative control: Kim and Pass] Jniform density bound

[Displacement functional: the thesis]
Weak compactness and Sousin space theory

[J. Ma, 2023]

Let $(M,d_{\rm g})$ be a complete Riemannian manifold with a lower Ricci curvature bound. If a measure $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to the set of a.c. probability measures, then its unique barycenter $\mu_{\mathbb{P}}$ is also a.c..

Improvements compared to the previous work [Eliminate technical assumption] a.c. with uniformly bounded density functions

[No obscure dependency on compactness] confirm the impact of lower Ricci curvature bound on the regularity of barycenters

 $\mathbb{P} = \sum_{i=1}^n \lambda_i \, \delta_{
u_i} \implies \mu_{\mathbb{P}}$ can be constructed.

However, when \mathbb{P} is not finitely supported, its barycenter $\mu_{\mathbb{P}}$ is only accessible via consistency: as a limit of constructable barycenters $\mu_{\mathbb{P}_j}$ with \mathbb{P}_j finitely supported.

[Quantitative control: Kim and Pass] Uniform density bound

[Displacement functional: the thesis]

weak compactness and Sousin space theory

[J. Ma, 2023]

Let $(M,d_{\rm g})$ be a complete Riemannian manifold with a lower Ricci curvature bound. If a measure $\mathbb{P}\in\mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to the set of a.c. probability measures, then its unique barycenter $\mu_{\mathbb{P}}$ is also a.c..

Improvements compared to the previous work
[Eliminate technical assumption]
a.c. with uniformly bounded density functions

[No obscure dependency on compactness] confirm the impact of lower Ricci curvature bound on the regularity of barycenters

```
\mathbb{P} = \sum_{i=1}^n \lambda_i \, \delta_{\nu_i} \implies \mu_{\mathbb{P}} \text{ can be constructed.} However, when \mathbb{P} is not finitely supported, its barycenter \mu_{\mathbb{P}} is only accessible via consistency: as a limit of constructable barycenters \mu_{\mathbb{P}_j} with \mathbb{P}_j finitely supported.
```

Quantitative control: Kim and Pass]
Uniform density bound
Displacement functional: the thesis]

[J. Ma, 2023]

Let (M,d_g) be a complete Riemannian manifold with a lower Ricci curvature bound. If a measure $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to the set of a.c. probability measures, then its unique barycenter $\mu_{\mathbb{P}}$ is also a.c..

Improvements compared to the previous work
[Eliminate technical assumption]
a.c. with uniformly bounded density functions

[No obscure dependency on compactness

[No obscure dependency on compactness] confirm the impact of lower Ricci curvature bound on the regularity of barycenters

Difficulty: control $\mu_{\mathbb{P}}$ via approximation

$$\mathbb{P} = \sum_{i=1}^n \lambda_i \, \delta_{
u_i} \implies \mu_{\mathbb{P}}$$
 can be constructed.

However, when $\mathbb P$ is not finitely supported, its barycenter $\mu_{\mathbb P}$ is only accessible via consistency: as a limit of constructable barycenters $\mu_{\mathbb P_j}$ with $\mathbb P_j$ finitely supported.

[Quantitative control: Kim and Pass] Uniform density bound

[Displacement functional: the thesis] Weak compactness and Souslin space theory

[J. Ma, 2023]

Let (M,d_g) be a complete Riemannian manifold with a lower Ricci curvature bound. If a measure $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to the set of a.c. probability measures, then its unique barycenter $\mu_{\mathbb{P}}$ is also a.c..

Improvements compared to the previous work
[Eliminate technical assumption]
a.c. with uniformly bounded density functions

[No obscure dependency on compactness] confirm the impact of lower Ricci curvature bound on the regularity of barycenters

Difficulty: control $\mu_{\mathbb{P}}$ via approximation

$$\mathbb{P} = \sum_{i=1}^n \lambda_i \, \delta_{
u_i} \implies \mu_{\mathbb{P}}$$
 can be constructed.

However, when $\mathbb P$ is not finitely supported, its barycenter $\mu_{\mathbb P}$ is only accessible via consistency: as a limit of constructable barycenters $\mu_{\mathbb P_j}$ with $\mathbb P_j$ finitely supported.

[Quantitative control: Kim and Pass] Uniform density bound

[Displacement functional: the thesis] Weak compactness and Souslin space theory

[J. Ma, 2023]

Let (M,d_g) be a complete Riemannian manifold with a lower Ricci curvature bound. If a measure $\mathbb{P} \in \mathcal{W}_2(\mathcal{W}_2(M))$ assigns positive mass to the set of a.c. probability measures, then its unique barycenter $\mu_{\mathbb{P}}$ is also a.c..

Improvements compared to the previous work
[Eliminate technical assumption]
a.c. with uniformly bounded density functions

[No obscure dependency on compactness] confirm the impact of lower Ricci curvature bound on the regularity of barycenters

Difficulty: control $\mu_{\mathbb{P}}$ via approximation

$$\mathbb{P} = \sum_{i=1}^n \lambda_i \, \delta_{
u_i} \implies \mu_{\mathbb{P}}$$
 can be constructed.

However, when $\mathbb P$ is not finitely supported, its barycenter $\mu_{\mathbb P}$ is only accessible via consistency: as a limit of constructable barycenters $\mu_{\mathbb P_j}$ with $\mathbb P_j$ finitely supported.

[Quantitative control: Kim and Pass] Uniform density bound

[Displacement functional: the thesis] Weak compactness and Souslin space theory

Displacement functionals defined for a.c. measures

 $[(M, d_g)]$: m-dimensional, with lower Ricci curvature bound 0 (for simplicity).

 $[\mathbb{P}]: \mathbb{P} = \sum_{j=1}^n \lambda_j \, \delta_{\nu_j}$, each $\mathrm{supp}(\nu_j)$ is compact, $\{\nu_i\}_{1 \leq i \leq k}$ are a.c., $\Lambda := \sum_{i=1}^k \lambda_i$.

[G]: $\mathcal{G}(f \cdot \mathrm{Vol}) = \int_M G(f) \, \mathrm{d} \, \mathrm{Vol}$, G satisfies certain properties (defining $L_G > 0$).

$$\mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^{k} \frac{\lambda_i}{\Lambda} \mathcal{G}(\nu_i) + \frac{L_G}{2\Lambda} (m^2 + 2m).$$

Consider the entropy functional Ent with $G(x) := x \log x$ and n = k = 2

$$\operatorname{Ent}(\mu_{\mathbb{P}}) \le \lambda_1 \operatorname{Ent}(\nu_1) + \lambda_2 \operatorname{Ent}(\nu_2)$$

Displacement functionals defined for a.c. measures

 $[(M, d_q)]$: m-dimensional, with lower Ricci curvature bound 0 (for simplicity).

 $[\mathbb{P}]: \mathbb{P} = \sum_{j=1}^n \lambda_j \, \delta_{\nu_j}$, each $\operatorname{supp}(\nu_j)$ is compact, $\{\nu_i\}_{1 \leq i \leq k}$ are a.c., $\Lambda := \sum_{i=1}^k \lambda_i$.

 $[\mathcal{G}]$: $\mathcal{G}(f \cdot \text{Vol}) = \int_M G(f) \, d \, \text{Vol}, \, G$ satisfies certain properties (defining $L_G > 0$).

$$\mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^k rac{\lambda_i}{\Lambda} \mathcal{G}(\nu_i) + rac{L_G}{2\Lambda} (m^2 + 2m).$$

Background: synthetic lower Ricci curvature bound 0

Consider the entropy functional Ent with $G(x) := x \log x$ and n = k = 2. (M, d_q) has lower Ricci curvature bound 0 iff:

$$\operatorname{Ent}(\mu_{\mathbb{P}}) \leq \lambda_1 \operatorname{Ent}(\nu_1) + \lambda_2 \operatorname{Ent}(\nu_2)$$

Auxiliary sets $B(G, L) := \{ \nu \mid \mathcal{G}(\nu) \leq L \}$

According to the previous inequality

$$\begin{split} \mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^k \frac{\lambda_i}{\Lambda} \mathcal{G}(\nu_i) + \frac{\underline{L_G}}{2\Lambda} (m^2 + 2m), \\ \text{if } \{\nu_i\}_{1 \leq i \leq k} \subset \mathrm{B}(G,L) \text{, then } \mu_{\mathbb{P}} \in \mathrm{B}(G,L'). \end{split}$$

Properties imposed on G

- a. function $H(x) := e^{-x} G(e^x)$ has positive and continuous derivative bounded from above by L_G
- b. continuous, convex, positive with G(0) = 0

c.
$$\lim_{x \to \infty} \frac{G(x)}{x} = +\infty$$

d. increasing

Auxiliary sets $B(G, L) := \{ \nu \mid \mathcal{G}(\nu) \leq L \}$

According to the previous inequality

thus is a c

$$\mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^k rac{\lambda_i}{\Lambda} \mathcal{G}(
u_i) + rac{L_G}{2\Lambda}(m^2 + 2m),$$

 $\inf \ \{\nu_i\}_{1 \leq i \leq k} \subset \mathrm{B}(G,L) \text{, then } \mu_{\mathbb{P}} \in \mathrm{B}(G,L').$

Properties imposed on ${\it G}$

- a. function $H(x):=e^{-x}\;G(e^x)$ has positive and continuous derivative bounded from above by L_G
- b. continuous, convex, positive with ${\it G}(0)=0$

c.
$$\lim_{x \to \infty} \frac{G(x)}{x} = +\infty$$

d. increasing

Bound $\mathcal G$ in approximation Compactness in $\sigma(L^1,L^\infty)$ Souslin space theory $\downarrow \hspace{1cm} \downarrow \hspace{$

to a B(G, L) set.

w.r.t. $\sigma(L^1, L^{\infty})$.

Auxiliary sets $B(G, L) := \{ \nu \mid \mathcal{G}(\nu) \leq L \}$

According to the previous inequality

$$\mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^k \frac{\lambda_i}{\Lambda} \mathcal{G}(\nu_i) + \frac{L_G}{2\Lambda}(m^2 + 2m),$$
 if $\{\nu_i\}_{1 \leq i \leq k} \subset \mathrm{B}(G,L)$, then $\mu_{\mathbb{P}} \in \mathrm{B}(G,L')$.

Properties imposed on G

- a. function $H(x) := e^{-x} G(e^x)$ has positive and continuous derivative bounded from above by L_G
- b. continuous, convex, positive with G(0) = 0

c.
$$\lim_{x \to \infty} \frac{G(x)}{x} = +\infty$$

d. increasing

Bound \mathcal{G} in approximation

Compactness in $\sigma(L^1, L^{\infty})$

Souslin space theory

1. G is lower semi-continuous:

$$\mathcal{G}(\lim_{l\to+\infty}\mu_l)\leq \liminf_{l\to+\infty}\mathcal{G}(\mu_l).$$

$$\mathbb{P}_{j}(\mathsf{B}(G,L+1)) \geq \mathbb{P}(\mathsf{B}(G,L)) > 0.$$

Auxiliary sets $B(G, L) := \{ \nu \mid \mathcal{G}(\nu) \leq L \}$

According to the previous inequality

$$\mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^k \frac{\lambda_i}{\Lambda} \mathcal{G}(\nu_i) + \frac{L_G}{2\Lambda}(m^2 + 2m),$$
 if $\{\nu_i\}_{1 \leq i \leq k} \subset \mathsf{B}(G,L)$, then $\mu_{\mathbb{P}} \in \mathsf{B}(G,L')$.

Properties imposed on G

- a. function $H(x) := e^{-x} G(e^x)$ has positive and continuous derivative bounded from above by $L_{\it C}$
- b. continuous, convex, positive with G(0) = 0

c.
$$\lim_{x \to \infty} \frac{G(x)}{x} = +\infty$$

d. increasing

Bound \mathcal{G} in approximation

Compactness in $\sigma(L^1, L^{\infty})$

Souslin space theory

1. G is lower semi-continuous:

$$\mathcal{G}(\lim_{l\to+\infty}\mu_l)\leq \liminf_{l\to+\infty}\mathcal{G}(\mu_l).$$

2. For $\mathbb{P}_i \to \mathbb{P}$ and large i.

$$\mathbb{P}_{j}(\mathsf{B}(G,L+1)) \geq \mathbb{P}(\mathsf{B}(G,L)) > 0.$$

Auxiliary sets $B(G, L) := \{ \nu \mid \mathcal{G}(\nu) \leq L \}$

According to the previous inequality

$$\mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^k rac{\lambda_i}{\Lambda} \mathcal{G}(
u_i) + rac{L_G}{2\Lambda}(m^2 + 2m),$$

 $\text{if }\{\nu_i\}_{1\leq i\leq k}\subset \mathrm{B}(G,L)\text{, then }\mu_{\mathbb{P}}\in \mathrm{B}(G,L').$

Properties imposed on ${\it G}$

- a. function $H(x):=e^{-x}\;G(e^x)$ has positive and continuous derivative bounded from above by L_G
- b. continuous, convex, positive with ${\it G}(0)=0$

c.
$$\lim_{x \to \infty} \frac{G(x)}{x} = +\infty$$

d. increasing

Bound $\mathcal G$ in approximation

1. \mathcal{G} is lower semi-continuous:

$$\mathcal{G}(\lim_{l\to+\infty}\mu_l)\leq \liminf_{l\to+\infty}\mathcal{G}(\mu_l).$$

2. For $\mathbb{P}_j \to \mathbb{P}$ and large j,

$$\mathbb{P}_{j}(\mathsf{B}(G,L+1)) \geq \mathbb{P}(\mathsf{B}(G,L)) > 0.$$

Compactness in $\sigma(L^1, L^{\infty})$

Dunford-Pettis theorem

de la Vallée Poussin criterion

If F is weakly compact, then it is uniformly integrable, and thus contained in a B(G,L) set.

Souslin space theory

Auxiliary sets $B(G, L) := \{ \nu \mid \mathcal{G}(\nu) \leq L \}$

According to the previous inequality

$$\mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^k rac{\lambda_i}{\Lambda} \mathcal{G}(
u_i) + rac{L_G}{2\Lambda}(m^2 + 2m),$$

 $\inf \ \{\nu_i\}_{1 \leq i \leq k} \subset \mathrm{B}(G,L) \text{, then } \mu_{\mathbb{P}} \in \mathrm{B}(G,L').$

Properties imposed on ${\it G}$

- a. function $H(x):=e^{-x}\,G(e^x)$ has positive and continuous derivative bounded from above by L_G
- b. continuous, convex, positive with G(0) = 0

c.
$$\lim_{x \to \infty} \frac{G(x)}{x} = +\infty$$

d. increasing

Bound $\mathcal G$ in approximation

1. \mathcal{G} is lower semi-continuous:

$$\mathcal{G}(\lim_{l\to+\infty}\mu_l)\leq \liminf_{l\to+\infty}\mathcal{G}(\mu_l).$$

2. For $\mathbb{P}_j \to \mathbb{P}$ and large j,

$$\mathbb{P}_{j}(\mathsf{B}(G,L+1)) \geq \mathbb{P}(\mathsf{B}(G,L)) > 0.$$

Compactness in $\sigma(L^1, L^{\infty})$

Dunford-Pettis theorem de la Vallée Poussin criterion

If F is weakly compact, then it is uniformly integrable, and thus contained in a $\mathrm{B}(G,L)$ set.

Souslin space theory

Auxiliary sets $B(G, L) := \{ \nu \mid \mathcal{G}(\nu) \leq L \}$

According to the previous inequality

$$\mathcal{G}(\mu_{\mathbb{P}}) \leq \sum_{i=1}^{k} \frac{\lambda_i}{\Lambda} \mathcal{G}(\nu_i) + \frac{L_G}{2\Lambda} (m^2 + 2m),$$

if $\{\nu_i\}_{1\leq i\leq k}\subset \mathrm{B}(G,L)$, then $\mu_{\mathbb{P}}\in \mathrm{B}(G,L')$.

Properties imposed on ${\it G}$

- a. function $H(x):=e^{-x}\;G(e^x)$ has positive and continuous derivative bounded from above by L_G
- b. continuous, convex, positive with G(0)=0

c.
$$\lim_{x \to \infty} \frac{G(x)}{x} = +\infty$$

d. increasing

Bound $\mathcal G$ in approximation

1. \mathcal{G} is lower semi-continuous:

$$\mathcal{G}(\lim_{l\to+\infty}\mu_l)\leq \liminf_{l\to+\infty}\mathcal{G}(\mu_l).$$

2. For $\mathbb{P}_j \to \mathbb{P}$ and large j,

$$\mathbb{P}_{j}(\mathsf{B}(G,L+1)) \geq \mathbb{P}(\mathsf{B}(G,L)) > 0.$$

Compactness in $\sigma(L^1, L^{\infty})$

de la Vallée Poussin criterion


Dunford-Pettis theorem

If F is weakly compact, then it is uniformly integrable, and thus contained in a $\mathrm{B}(G,L)$ set.


Souslin space theory

 $\mathbb P$ is a Radon measure w.r.t. the $\sigma(L^1,L^\infty)$ topology.


Shifting focus: manifolds \longrightarrow metric trees



Shifting focus: manifolds \longrightarrow metric trees



Reduction Technique: flatten the tree to solve optimal transport problems




[Metric tree $\Gamma = (\mathcal{V}, \mathcal{E}, d_l)$]

Metric d_l is induced by length function $l: \mathcal{E} \to \mathbb{R}$.


 $\boldsymbol{\Gamma}$ is a proper and geodesic metric space.

Reduction map $T^{ec{e}}:\Gamma ightarrow \mathbb{R}$ associated to $ec{e}$



```
[Flatten via the reduction map T^{\vec{e}}] T^{\vec{e}} is continuous. T^{\vec{e}}(\vec{e}) = [0, l(\vec{e})]. For x \in \vec{e}, y \in \Gamma, d_l(x,y) = |T^{\vec{e}}(x) - T^{\vec{e}}(y)|.
```

Reduction map $T^{ec{e}}:\Gamma ightarrow \mathbb{R}$ associated to $ec{e}$



[Flatten via the reduction map $T^{\vec{e}}$] $T^{\vec{e}}$ is continuous. $T^{\vec{e}}(\vec{e}) = [0, l(\vec{e})]$. For $x \in \vec{e}, y \in \Gamma$, $d_l(x,y) = |T^{\vec{e}}(x) - T^{\vec{e}}(y)|$.


Reduction preserves Wasserstein distances

For $\mu, \nu \in \mathcal{W}_2(\Gamma)$, if $\operatorname{supp}(\mu) \subset \vec{e}$, then

$$d_W(\mu,\nu) = d_W(\mathcal{T}(\mu), \mathcal{T}(\nu)),$$

where $\mathcal{T}: \mathcal{W}_2(\Gamma) \to \mathcal{W}_2(\mathbb{R})$ is the induced push-forward map: $\mathcal{T}(\mu) := T^{\vec{e}}_{\#}\mu$.

Reduction map $T^{ec{e}}:\Gamma ightarrow \mathbb{R}$ associated to $ec{e}$



[Flatten via the reduction map $T^{\vec{e}}$] $T^{\vec{e}}$ is continuous. $T^{\vec{e}}(\vec{e}) = [0, l(\vec{e})]$. For $x \in \vec{e}, y \in \Gamma$, $d_l(x,y) = |T^{\vec{e}}(x) - T^{\vec{e}}(y)|$.

Reduction preserves Wasserstein distances


For $\mu, \nu \in \mathcal{W}_2(\Gamma)$, if $\operatorname{supp}(\mu) \subset \vec{e}$, then

$$d_W(\mu, \nu) = d_W(\mathcal{T}(\mu), \mathcal{T}(\nu)),$$

where $\mathcal{T}:\mathcal{W}_2(\Gamma) \to \mathcal{W}_2(\mathbb{R})$ is the induced push-forward map: $\mathcal{T}(\mu) := T^{\vec{e}}{}_\#\mu$.

Idea of its proof: recover optimal transports between from $\mathbb R$ to Γ via $\nu = \sum_{e \in \mathcal E} \lambda_e \nu_e$.

Reduction map $T^{ec{e}}:\Gamma ightarrow \mathbb{R}$ associated to $ec{e}$



[Flatten via the reduction map $T^{\vec{e}}$] $T^{\vec{e}}$ is continuous. $T^{\vec{e}}(\vec{e}) = [0, l(\vec{e})]$. For $x \in \vec{e}, y \in \Gamma$, $d_l(x, y) = |T^{\vec{e}}(x) - T^{\vec{e}}(y)|$.

Reduction preserves Wasserstein distances

For $\mu, \nu \in \mathcal{W}_2(\Gamma)$, if $\operatorname{supp}(\mu) \subset \vec{e}$, then

$$d_W(\mu,\nu) = d_W(\mathcal{T}(\mu), \mathcal{T}(\nu)),$$

where $\mathcal{T}:\mathcal{W}_2(\Gamma)\to\mathcal{W}_2(\mathbb{R})$ is the induced push-forward map: $\mathcal{T}(\mu):=T^{\vec{e}}_{\ \#}\mu$.

Idea of its proof: recover optimal transports between from $\mathbb R$ to Γ via $\nu = \sum_{e \in \mathcal E} \lambda_e \nu_e$.

Properties: 1. $\mathcal{T}(\nu)$ is a.c. $\Leftrightarrow \nu$ is a.c.; 2. if $\mathcal{T}(\nu)$ is singular, then so is ν .

Restriction property of Wasserstein barycenters

Assume the base metric space $\left(E,d\right)$ is proper. For any decomposition of a given barycenter

$$\mu_{\mathbb{P}} = \lambda \,\mu^1 + (1 - \lambda)\mu^2,$$

there exist continuous maps F^i such that

- 1. $\lambda F^{1}(\nu) + (1 \lambda)F^{2}(\nu) = \nu$;
- 2. μ^i is a barycenter of $\mathbb{Q}^i := F^i{}_\# \mathbb{P}$.

Restriction property of Wasserstein barycenters

Assume the base metric space $\left(E,d\right)$ is proper. For any decomposition of a given barycenter

$$\mu_{\mathbb{P}} = \lambda \,\mu^1 + (1 - \lambda)\mu^2,$$

there exist continuous maps F^i such that

- 1. $\lambda F^{1}(\nu) + (1 \lambda)F^{2}(\nu) = \nu$;
- 2. μ^i is a barycenter of $\mathbb{Q}^i := F^i_\# \mathbb{P}$.

[Preserved properties of \mathbb{P}]

For any reference measure η on E, if $\mathbb P$ assigns full (or positive) mass to measures a.c. w.r.t. η , then so do the measures $\mathbb Q^1$ and $\mathbb Q^2$.

Restriction property of Wasserstein barycenters

Assume the base metric space $\left(E,d\right)$ is proper. For any decomposition of a given barycenter

$$\mu_{\mathbb{P}} = \lambda \,\mu^1 + (1 - \lambda)\mu^2,$$

there exist continuous maps F^i such that

- 1. $\lambda F^{1}(\nu) + (1 \lambda)F^{2}(\nu) = \nu$;
- 2. μ^i is a barycenter of $\mathbb{Q}^i := F^i_\# \mathbb{P}$.

A lemma using reduction technique

Fix an oriented edge \vec{e} of a metric tree Γ . If a barycenter $\mu_{\mathbb{P}}(\mathring{e})=1$, then $\mathcal{T}(\mu_{\mathbb{P}})$ is the unique barycenter of $\mathcal{T}_{\#}\mathbb{P}$.

Restriction property of Wasserstein barycenters

Assume the base metric space $({\cal E},d)$ is proper. For any decomposition of a given barycenter

$$\mu_{\mathbb{P}} = \lambda \,\mu^1 + (1 - \lambda)\mu^2,$$

there exist continuous maps F^{i} such that

- 1. $\lambda F^{1}(\nu) + (1 \lambda)F^{2}(\nu) = \nu$;
- 2. μ^i is a barycenter of $\mathbb{Q}^i := F^i{}_\# \mathbb{P}$.

A lemma using reduction technique

Fix an oriented edge \vec{e} of a metric tree Γ . If a barycenter $\mu_{\mathbb{P}}(\mathring{e})=1$, then $\mathcal{T}(\mu_{\mathbb{P}})$ is the unique barycenter of $\mathcal{T}_{\#}\mathbb{P}$.

Application: almost absolute continuity on Γ If $\mathbb{P}(\text{a.c. measures}) > 0$, then $\mu_{\mathbb{P}}|_{\mathring{e}}$ is a.c. for any barycenter $\mu_{\mathbb{P}}$ and any edge \overrightarrow{e} .

Therefore, if $\mu_{\mathbb{P}}$ is not a.c., then its singular part is a sum of Dirac measures at vertices.

Rigid properties of barycenters on \mathbb{R}

A measure property $\mathcal Q$ is \emph{rigid} if

 $\mu_{\mathbb{P}} \text{ satisfies } \mathcal{Q} \implies \mathbb{P}\text{-almost every } \nu \text{ satisfies } \mathcal{Q}.$

Rigid properties of barycenters on $\mathbb R$

A measure property Q is *rigid* if

 $\mu_{\mathbb{P}} \text{ satisfies } \mathcal{Q} \implies \mathbb{P}\text{-almost every } \nu \text{ satisfies } \mathcal{Q}.$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property $\mathcal Q$ is rigid if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every ν satisfies $\mathcal{Q}.$

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property $\mathcal Q$ is rigid if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every ν satisfies $\mathcal{Q}.$

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

1. For 0 < t < 1, f_{μ}^{-1} is defined by

$$f_{\mu}^{-1}(t) := \inf_{x} \{ x \in \mathbb{R} \mid f_{\mu}(x) > t \}.$$

2. $f_{\mu}^{-1}(0)$, $f_{\mu}^{-1}(1)$: defined as one-sided limits.

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property $\mathcal Q$ is \emph{rigid} if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every ν satisfies $\mathcal{Q}.$

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) d \mathbb{P}(\nu)$$

[Quantile functions]

$$f_{\mu}^{-1}(0) = \inf \operatorname{supp}(\mu)$$

$$f_{\mu}^{-1}(1) = \sup \operatorname{supp}(\mu)$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property $\mathcal Q$ is \emph{rigid} if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every ν satisfies $\mathcal{Q}.$

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

$$\mu = \delta_x \Longleftrightarrow f_{\mu}^{-1} \equiv x$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property $\mathcal Q$ is \emph{rigid} if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every u satisfies \mathcal{Q} .

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) d \mathbb{P}(\nu)$$

[Quantile functions]

Consider the special case $supp(\mu) \subset [0,1]$.

 f_{μ} is singular $\iff f_{\mu}^{-1}$ is singular

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

Rigid properties of barycenters on $\mathbb R$

A measure property $\mathcal Q$ is rigid if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every ν satisfies $\mathcal{Q}.$

Explicit formula for barycenters


$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

Consider the special case $\operatorname{supp}(\mu) \subset [0,1].$

 f_{μ} is singular $\iff f_{\mu}^{-1}$ is singular

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set



$$ArcLength(f_{\mu}) = || Leb^{1} |_{[0,1]} + i \mu ||_{TV} = 2$$

Rigid properties of barycenters on $\mathbb R$

A measure property Q is *rigid* if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every ν satisfies $\mathcal{Q}.$

Explicit formula for barycenters


$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

Consider the special case $\operatorname{supp}(\mu) \subset [0,1]$.

 f_{μ} is singular $\iff f_{\mu}^{-1}$ is singular

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set



 $ArcLength(f_{\mu}) = 2 = ArcLength(f_{\mu}^{-1})$

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property $\mathcal Q$ is \emph{rigid} if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every u satisfies \mathcal{Q} .

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

 f_{μ}^{-1} is singular iff

$$V_0^1 (f_\mu^{-1} - \mathrm{Id}) = V_0^1 (f_\mu^{-1} + \mathrm{Id}).$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

$$V_0^1\left(f_{\mu_{\mathbb{P}}}^{-1}-\operatorname{Id}\right) \leq \int_{\mathcal{W}_2(\mathbb{R})} \, V_0^1\left(f_{\nu}^{-1}-\operatorname{Id}\right) \operatorname{d}\mathbb{P}(\nu)$$

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property $\mathcal Q$ is rigid if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every ν satisfies \mathcal{Q} .

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

 f_{μ}^{-1} is singular iff

$$V_0^1 (f_\mu^{-1} - \mathrm{Id}) = V_0^1 (f_\mu^{-1} + \mathrm{Id}).$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

$$\begin{split} V_0^1 \left(f_{\mu_{\mathbb{P}}}^{-1} - \operatorname{Id} \right) &\leq \int_{\mathcal{W}_2(\mathbb{R})} V_0^1 \left(f_{\nu}^{-1} - \operatorname{Id} \right) d \, \mathbb{P}(\nu) \\ &\leq \int_{\mathcal{W}_2(\mathbb{R})} V_0^1 \left(f_{\nu}^{-1} \right) + V_0^1 \left(\operatorname{Id} \right) d \, \mathbb{P}(\nu) \end{split}$$

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property Q is *rigid* if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every u satisfies \mathcal{Q} .

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

 f_{μ}^{-1} is singular iff

$$V_0^1 (f_\mu^{-1} - \mathrm{Id}) = V_0^1 (f_\mu^{-1} + \mathrm{Id}).$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

$$\begin{split} V_0^1 \left(f_{\mu_{\mathbb{P}}}^{-1} - \operatorname{Id} \right) &\leq \int_{\mathcal{W}_2(\mathbb{R})} V_0^1 \left(f_{\nu}^{-1} - \operatorname{Id} \right) d \, \mathbb{P}(\nu) \\ &\leq \int_{\mathcal{W}_2(\mathbb{R})} V_0^1 \left(f_{\nu}^{-1} \right) + V_0^1 \left(\operatorname{Id} \right) d \, \mathbb{P}(\nu) \\ &= \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(1) - f_{\nu}^{-1}(0) + 1 \, d \, \mathbb{P}(\nu) \end{split}$$

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property Q is rigid if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every u satisfies \mathcal{Q} .

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

 f_{μ}^{-1} is singular iff

$$V_0^1 (f_\mu^{-1} - \mathrm{Id}) = V_0^1 (f_\mu^{-1} + \mathrm{Id}).$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

$$\begin{split} V_0^1 \left(f_{\mu_{\mathbb{P}}}^{-1} - \mathrm{Id} \right) &\leq \int_{\mathcal{W}_2(\mathbb{R})} V_0^1 \left(f_{\nu}^{-1} - \mathrm{Id} \right) \mathrm{d} \, \mathbb{P}(\nu) \\ &\leq \int_{\mathcal{W}_2(\mathbb{R})} V_0^1 \left(f_{\nu}^{-1} \right) + V_0^1 \left(\mathrm{Id} \right) \mathrm{d} \, \mathbb{P}(\nu) \\ &= \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(1) - f_{\nu}^{-1}(0) + 1 \, \mathrm{d} \, \mathbb{P}(\nu) \\ &= f_{\mu_{\mathbb{P}}}^{-1}(1) - f_{\mu_{\mathbb{P}}}^{-1}(0) + 1 \end{split}$$

Rigid properties of barycenters on $\ensuremath{\mathbb{R}}$

A measure property Q is rigid if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every u satisfies \mathcal{Q} .

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) \, \mathrm{d} \, \mathbb{P}(\nu)$$

[Quantile functions]

 f_{μ}^{-1} is singular iff

$$V_0^1 (f_\mu^{-1} - \mathrm{Id}) = V_0^1 (f_\mu^{-1} + \mathrm{Id}).$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set

$$\begin{split} V_0^1 \left(f_{\mu_{\mathbb{P}}}^{-1} - \operatorname{Id} \right) &\leq \int_{\mathcal{W}_2(\mathbb{R})} V_0^1 \left(f_{\nu}^{-1} - \operatorname{Id} \right) \operatorname{d} \mathbb{P}(\nu) \\ &\leq \int_{\mathcal{W}_2(\mathbb{R})} V_0^1 \left(f_{\nu}^{-1} \right) + V_0^1 \left(\operatorname{Id} \right) \operatorname{d} \mathbb{P}(\nu) \\ &= \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(1) - f_{\nu}^{-1}(0) + 1 \operatorname{d} \mathbb{P}(\nu) \\ &= f_{\mu_{\mathbb{P}}}^{-1}(1) - f_{\mu_{\mathbb{P}}}^{-1}(0) + 1 \\ &= V_0^1 \left(f_{\mu_{\mathbb{P}}}^{-1} + \operatorname{Id} \right) \end{split}$$

Rigid properties of barycenters on $\mathbb R$

A measure property Q is *rigid* if

 $\mu_{\mathbb{P}}$ satisfies $\mathcal{Q} \implies \mathbb{P}$ -almost every ν satisfies $\mathcal{Q}.$

Explicit formula for barycenters

$$f_{\mu_{\mathbb{P}}}^{-1}(t) = \int_{\mathcal{W}_2(\mathbb{R})} f_{\nu}^{-1}(t) d \mathbb{P}(\nu)$$

- a. being not a.c.
- b. having compact support
- c. being a Dirac measure
- d. being singular
- e. being supported in a negligible set [skip]

Future research directions

- 1. Extending displacement functional arguments to metric measure spaces
- 2. Quantitative estimates for barycenter densities
- 3. Necessity of curvature bounds and the role of branching
- 4. Generalizing reduction techniques to metric graphs

Example: non-uniqueness of Wasserstein barycenters on metric trees

